arXiv:1304.3045v1 [physics.atom-ph] 10 Apr 2013

Do we have a new capture mechanism?

Yu. V. Popov*
Skobeltsyn Institute of Nuclear Physics,
Lomonosov Moscow State University, Moscow 119991, Russia

V. L. Shablov
Obninsk Institute for Nuclear Power Engineering,
National Research Nuclear University,
Obninsk, Kaluga region 249040, Russia

K. A. Kouzakov
Faculty of Physics, Lomonosov Moscow State University, Moscow 119991, Russia and
Skobeltsyn Institute of Nuclear Physics,
Lomonosov Moscow State University, Moscow 119991, Russia

A. G. Galstyan
Faculty of Physics, Lomonosov Moscow State University, Moscow 119991, Russia

Abstract

We inspect the first-order electron-electron capture scenario for transfer ionization that has been recently formulated by Voitkiv *et al.* (Phys. Rev. A **86**, 012709 (2012) and references therein). Using the multichannel scattering theory for many-body systems with Coulomb interactions, we show that this scenario is just a part of the well-studied Oppenheimer-Brinkmann-Kramers approximation. Accurate numerical calculations for the proton-helium transfer ionization reaction in the first Born approximation exhibit no appreciable manifestation of the claimed mechanism.

PACS numbers: 34.70.+e, 34.10.+x, 34.50.Fa

---

*Electronic address: popov@srd.sinp.msu.ru

I. INTRODUCTION

Various capture processes, including single and double charge transfer, transfer excitation (TE), and transfer ionization (TI), have been attracting interest from researchers for nearly a hundred years. Recent $H^+ + He$ experiments allow to distinguish not only the ground state of the residual ion He^+, but also its excited and ionization states, as well as the ground and excited states of the final atomic hydrogen [1–4]. To our knowledge, the first (classical) description of the capture mechanism was given by Thomas in 1927 [5]. Briefly, an electron, after being kicked by a fast proton, elastically scatters from the target nucleus. The scattered electron can move parallel to the proton with the same velocity. This is a classical description of the capture process. To transfer the electron into a hydrogen bound state, a gamma-quant must be emitted to carry away the released energy (the so-called radiative capture [6]). This mechanism is referred to as Ne-Thomas.

Almost at the same time, a quantum mechanical explanation how a target electron can be captured into a bound state of a fast moving projectile (proton), was given by Oppenheimer, Brinkmann and Kramers (OBK) [7]. In the OBK scenario the electron transfer proceeds via an overlap of initial and final wave functions of the projectile-target system. This so-called kinematical capture relies strongly on the radial and angular electron correlations in the target, if we consider TE and TI processes. It must be noted that in quantum mechanics the transition of an electron to the hydrogen bound state can be nonradiative, and the energy excess can be carried away by a third body that participates in the reaction. In 1974, Shakeshaft [8] demonstrated connection between the correct classical and quantum mechanical second-order treatments of the Ne-Thomas process at high projectile velocities. And later, Briggs and Taulbjerg [9] carried out a quantum treatment of the so-called ee-Thomas mechanism, where an electron after the first kick is scattered by another target electron, not by the nucleus.

In quantum mechanics fast processes are usually treated in terms of Born approximations. This framework is directly applicable in the case of two-body scattering, but requires additional careful considerations in the many-body case. The situation of particular importance is when the entrance channel of the reaction is different from its exit channel, for example, as it is in capture processes. Within the multichannel scattering theory, the OBK mechanism can be attributed to the first Born approximation (FBA), whereas the Ne- and *ee*-Thomas mechanisms can be described using the second Born approximation (SBA). Any Born approximation is a sum of matrix elements. Each of them corresponds to a particular interaction that enters a total perturbation potential. For example, the OBK matrix element is one of the three FBA terms. The SBA contains twelve different terms, and only two of them correspond to the *Ne*- and *ee*-Thomas mechanisms.

Recently Voitkiv *et al.* published a series of papers [10–12] putting forth a new first-order capture mechanism that can be called electron-electron Auger (or *ee*-Auger) [12]. According to this mechanism, the electron undergoes a nonradiative transition from the atomic state to the bound state of the projectile, transferring the energy excess to the another atomic electron which is emitted from the atom. This scenario resembles a kind of Auger decay and to be contrasted with the first-order radiative capture [13] which is accompanied by emission of a photon instead of an electron. A clear signature of the *ee*-Auger mechanism, according to Voitkiv *et al.*, is emission of the electron in the direction opposite to the projectile motion (in the rest frame of the atom). In this work, we examine the mechanism suggested by Voitkiv *et al.* on the basis of consistent multichannel scattering theory. We show that, in contrast to the claim of Ref. [10], it is not new and previously undiscussed. Namely, it is just a part of the usual kinematic capture in the OBK approximation, and the correlated electron-electron emission is nothing else but a result of the electron-electron correlations in the target atom.

The paper is organized as follows. Sec. II is devoted to quantum scattering theory of the considered transfer ionization process. In Sec. III, effects of distortion beyond the first-order approximation are discussed. Numerical results are presented in Sec. IV. Finally, Sec. V summarizes this work. Atomic units (a.u., $\hbar = e = m_e = 1$) are used throughout unless otherwise specified.

II. MULTICHANNEL SCATTERING THEORY

In this section we remind basic formulas of quantum scattering theory for many-body systems. More mathematical details, particularly for the case involving charged fragments, one can find in the review article [14]. A set of relative momenta defining motion of n_α fragments colliding in the asymptotic channel is denoted by $\vec{p}_\alpha$. In turn, the ket vector $|\phi_\alpha\rangle$ stands for a product of bound (spectral) state wave functions, which define the channel

α. Hence, the ket vector $|\phi_\alpha, \vec{p}_\alpha\rangle$ is the eigenfunction of the asymptotic hamiltonian H_α: $(E_\alpha - H_\alpha)|\phi_\alpha, \vec{p}_\alpha\rangle = 0$. The total hamiltonian is $H = H_\alpha + V_\alpha$, where V_α is a set of two-body interaction potentials, which we consider as perturbation, and they define the terms of the Born series: FBA, SBA, and so on.

The amplitude of the transition from the channel α to the channel β can be presented using two forms. These are the *post*-form

$$T_{\beta\alpha}(E) = \langle\phi_\beta, \vec{p}_\beta|V_\beta + V_\beta(E - H + i0)^{-1}V_\alpha|\phi_\alpha, \vec{p}_\alpha\rangle, \tag{1}$$

and the *prior*-form

$$\tilde{T}_{\beta\alpha}(E) = \langle\phi_\beta, \vec{p}_\beta|V_\alpha + V_\beta(E - H + i0)^{-1}V_\alpha|\phi_\alpha, \vec{p}_\alpha\rangle. \tag{2}$$

It is straightforward to demonstrate that on the energy shell, where $E = E_\alpha = E_\beta$, one has (see, for instance, Ref. [15])

$$T_{\beta\alpha}(E) = \tilde{T}_{\beta\alpha}(E). \tag{3}$$

Moreover, since $V_\alpha = H - H_\alpha$ and $V_\beta = H - H_\beta$, the relation (3) holds true in the FBA case as well, that is, on the energy shell the FBA *post*- and *prior*-amplitudes coincide,

$$\langle\phi_\beta, \vec{p}_\beta|V_\beta|\phi_\alpha, \vec{p}_\alpha\rangle = \langle\phi_\beta, \vec{p}_\beta|V_\alpha|\phi_\alpha, \vec{p}_\alpha\rangle.$$

In the case of charged fragments the scattering theory should be modified. This modification is due to the fact that charged fragments continue to interact even at asymptotically large distances from each other, and their trajectories are not straight lines anymore. This asymptotic interaction in the channel α is characterized by the so-called Sommerfeld (Coulomb) parameter

$$\eta_\alpha = \sum_{i>j} \frac{Z_i Z_j}{v_{ij}}.$$

Here Z_i is the total charge of the ith asymptotic fragment (ion, atom, and electron), and v_{ij} the relative velocity of fragments i and j. At this point we introduce distorting potentials and distorted waves. There are many ways of doing this, but a channel distorted function $\chi_\alpha^\pm(\vec{p}_\alpha)$ must obey the asymptotic relation at $t \to \mp\infty$ [16]

$$e^{-iH_\alpha t}|\chi_\alpha^\pm(\vec{p}_\alpha)\rangle \to e^{-iE_\alpha t \pm i\eta_\alpha \ln|t| \pm iA_\alpha(\vec{p}_\alpha)}|\phi_\alpha, \vec{p}_\alpha\rangle. \tag{4}$$

Here $A_\alpha(\vec{p}_\alpha)$ is the Dollard phase, whose explicit form is not important for the following. A distorting potential (operator) must satisfy the only condition

$$\mathbb{V}_\alpha^\pm |\phi_\alpha, \vec{p}_\alpha\rangle = |\chi_\alpha^\pm(\vec{p}_\alpha)\rangle,$$

and the channel wave function $\Psi_\alpha^\pm(\vec{p}_\alpha)$ is given by the modified Lippmann-Schwinger representation [14, 17]

$$|\Psi_\alpha^\pm(\vec{p}_\alpha)\rangle = |\chi_\alpha^\pm(\vec{p}_\alpha)\rangle + (E_\alpha - H \pm i0)^{-1}(V_\alpha \mathbb{V}_\alpha^\pm + [H_\alpha, \mathbb{V}_\alpha^\pm])|\phi_\alpha, \vec{p}_\alpha\rangle. \tag{5}$$

All well known approximations (the Peterkop effective charges [18], CDW [19, 20], and etc.) are related to Eq. (5). The only difference between them is the choice of the operator $\mathbb{V}_\alpha^\pm$. Thus, we can present the *post*- and *prior*-amplitudes as ($\beta \neq \alpha,\ E = E_\alpha = E_\beta$) [14]

$$T_{\beta\alpha}(E) = \langle \phi_\beta, \vec{p}_\beta | (V_\beta \mathbb{V}_\beta^- + [H_\beta, \mathbb{V}_\beta^-])^+ | \Psi_\alpha^+(\vec{p}_\alpha)\rangle, \tag{6}$$

$$\tilde{T}_{\beta\alpha}(E) = \langle \Psi_\beta^-(\vec{p}_\beta) | V_\alpha \mathbb{V}_\alpha^+ + [H_\alpha, \mathbb{V}_\alpha^+] | \phi_\alpha, \vec{p}_\alpha\rangle. \tag{7}$$

These relations reduce to Eqs. (1) and (2), respectively, when $\mathbb{V}_{\alpha,\beta}^\pm = 1$ or, equivalently, $\eta_{\alpha,\beta} = 0$. As previously, the *post*- and *prior*-amplitudes coincide. Another important remark is that, if we account exactly for a particular distorting potential in the channel α, for example, for the Coulomb potential between two fragments, we must subtract the corresponding Sommerfeld parameter from the total channel parameter η_α (see, for instance, Ref. [14]).

Let us apply the above general formulas to the fast TI reaction $\mathrm{H}^+ + \mathrm{He} \to \mathrm{H} + \mathrm{He}^{2+} + e$ discussed in the papers of Voitkiv *et al.* [10–12]. The authors utilize the *post*-amplitude, which in the nonsymmetrized FBA can be written as

$$T_{fi}^{FBA}(E) = \langle \phi_{p1}, \varphi_{N2}^-(\vec{k}), \vec{p}_H | V_{N1} + V_{p2} + V_{12} + V_{Np} | \Phi_0, \vec{p}_0\rangle. \tag{8a}$$

In Eq. (8a), electrons are labelled by “1” and “2”, “p” labels the fast proton projectile, and “N” the target nucleus. The wave function $|\phi_{p1}\rangle$ is the bound (ground) state of atomic hydrogen, $|\varphi_{N2}(\vec{k})\rangle$ the continuum state of the He^+ ion, $|\Phi_0\rangle$ the helium wave function, $\vec{p}_0$ the proton momentum, $\vec{p}_H$ the hydrogen momentum, and $\vec{k}$ the momentum of the emitted electron. Further, Voitkiv *et al.* put forth their *ee*-Auger mechanism by leaving only the term V_{12} in the matrix element (8a). At the same time, they provide no evidence that this specific term indeed dominates in the FBA amplitude (8a). Moreover, taking into account that this amplitude is equal to that in the *prior*-form

$$\tilde{T}_{fi}^{FBA}(E) = \langle \phi_{p1}, \varphi_{N2}^{-}(\vec{k}), \vec{p}_H | V_{p1} + V_{p2} + V_{Np} | \Phi_0, \vec{p}_0 \rangle, \tag{8b}$$

there is no ground for such a claim. It was shown many times (see, for instance, Ref. [23]) that in the FBA *prior*-amplitude (8b) even the OBK term, corresponding to V_{p1} and containing the *ee*-Auger mechanism (see below), is not dominant in some kinematical situations. In other words, all four terms in (8a) must be taken into account. From the equality of the FBA amplitudes (8a) and (8b) we find that

$$\langle \phi_{p1}, \varphi_{N2}^{-}(\vec{k}), \vec{p}_H | V_{p1} | \Phi_0, \vec{p}_0 \rangle = \langle \phi_{p1}, \varphi_{N2}^{-}(\vec{k}), \vec{p}_H | V_{N1} + V_{12} | \Phi_0, \vec{p}_0 \rangle. \tag{9}$$

It follows that the OBK matrix element, occurring in the left-hand side, is equal to the sum of the two matrix elements generated by the potentials V_{N1} and V_{12}, respectively. Thus, it appears that the *ee*-Auger mechanism claimed by Voitkiv *et al.* as new and so far unconsidered is, in fact, already included in the very well studied OBK scenario.

Eqs. (8a) and (8b), are valid when the total Sommerfeld parameters in the initial and final channels, η_α and η_β, are equal to zero. In the present case, it is clearly fulfilled in the initial channel (the He atom is neutral and does not interact with the charged proton at asymptotically large separation distances). But it is also fulfilled in the final channel, because we use in (8) the spectral Coulomb functions $\varphi_{N2}^{-}(\vec{k})$ instead of plain waves and, as remarked above, the Sommerfeld parameter of the $N2$ subsystem does not contribute to the total one.

Now we consider the Schrödinger equation for the discussed problem in real space. Below we shortly recapitulate results from Ref. [2]. Let us introduce spatial positions of electrons in the laboratory frame $\vec{r}_1$, $\vec{r}_2$, and $\vec{R}$ is the position of the projectile proton. If we consider very small scattering angles (fractions of milliradians) and not too high projectile energies (within few MeV), then the target nucleus stays practically at rest in the laboratory frame during the collision and we can set its position as the origin of the coordinate system, i.e. $\vec{r}_N \approx 0$. We also introduce the position of electron 1 relative to the proton $\vec{x} = \vec{R} - \vec{r}_1$. After a simple paraxial transformation, $\Psi_f^{-}(\vec{p}_H, \vec{k}; \vec{R}, \vec{x}, \vec{r}_2) \to \exp(i\vec{p}_H \cdot \vec{R}) \Phi_f^{-}(\vec{k}; \vec{R}, \vec{x}, \vec{r}_2)$, we obtain for the wave function $|\Phi_f^-\rangle$ the following equation:

$$\left\{\left[\frac{1}{2(m+1)}\triangle_R + i\vec{v}_p\cdot\vec{\nabla}_R\right] + \left(\frac{1}{2}\triangle_x + \varepsilon_0^H - V_{p1}\right) + \left(\frac{1}{2}\triangle_2 + \frac{k^2}{2} - V_{N2}\right)\right\}\Phi_f^-(\vec{k};\vec{R},\vec{x},\vec{r}_2)$$
$$\approx (V_{p2} + V_{pN} + V_{N1} + V_{12})\ \Phi_f^-(\vec{k};\vec{R},\vec{x},\vec{r}_2), \tag{10}$$

with the final boundary condition $|\Phi_f^-\rangle \to |\phi_{p1},\varphi_{N2}^-(\vec{k})\rangle$. Since the proton mass is $m = 1836$, the term $\triangle_R/2(m+1)$ can be neglected in (10). Setting $R_z = v_p t$ and $\vec{R}_\perp = \vec{b}$, i.e., $\vec{R}(t) = \vec{v}_p t + \vec{b}$, we obtain the effective "time-dependent" Schrödinger equation

$$\left[i\frac{\partial}{\partial t} + \left(\frac{1}{2}\triangle_x + \varepsilon_0^H + \frac{1}{x}\right) + \left(\frac{1}{2}\triangle_2 + \frac{k^2}{2} + \frac{2}{r_2}\right)\right]\Phi_f^-(\vec{k};\vec{b},\vec{x},\vec{r}_2;t) =$$
$$\left(-\frac{1}{|\vec{R}(t)-\vec{r}_2|} + \frac{2}{R(t)} - \frac{2}{|\vec{R}(t)-\vec{x}|} + \frac{1}{|\vec{R}(t)-\vec{x}-\vec{r}_2|}\right)\Phi_f^-(\vec{k};\vec{b},\vec{x},\vec{r}_2;t). \tag{11}$$

In this connection, let us remind a well known relation between the present time-independent scattering theory and the approach of Refs. [10–12], where time-dependent formulation with classical treatment of the fast moving projectile is employed. In that approach, the heavy incoming ion (proton) is considered as a source of a classical field entering the time-dependent Schrödinger equation, and its trajectory in the laboratory frame is $\vec{R}(t) = \vec{b} + \vec{v}_p t,\quad (\vec{b}\cdot\vec{v}_p) = 0$, where b is the impact parameter. The time-dependent Schrödinger equation is

$$i\frac{\partial}{\partial t}|\Phi^\pm(t)\rangle = [H_0 + V_{p1}(t) + V_{p2}(t) + V_{Np}(t) + V_{N1} + V_{N2} + V_{12}]|\Phi^\pm(t)\rangle, \tag{12}$$

where H_0 is the kinetic-energy part of the total hamiltonian H. The interaction potential of the proton with the helium target is $V_{pHe}(t) = V_{p1}(t) + V_{p2}(t) + V_{Np}(t) \sim O(1/t^2)$, if $t \to \pm\infty$. The initial and final conditions at $t \to \mp\infty$, respectively, are

$$\langle\vec{r}_1,\vec{r}_2|\Phi^+(t\to-\infty)\rangle \to e^{-i\varepsilon_0^{He}t}\ \langle\vec{r}_1,\vec{r}_2|\Phi_0\rangle,$$

$$\langle\vec{x},\vec{r}_2|\Phi^-(t\to+\infty)\rangle \to e^{i(v_p^2/2-\varepsilon^H-k^2/2)t}\ e^{-i\vec{v}_p\cdot\vec{x}}\langle\vec{x},\vec{r}_2|\phi_{p1},\varphi_{N2}^-(\vec{k})\rangle,$$

where $\varepsilon_0^{He} \approx -2.9037$ is the helium ground-state energy, $\varepsilon^H = -0.5$ the ground energy of hydrogen atom, and $\vec{x} = \vec{R}(t) - \vec{r}_1$. The so-called translational factor $e^{i[(v_p^2/2)t - \vec{v}_p\cdot\vec{x}]}$ [9] arises as a result of Galilean transformation of the hydrogen state from the projectile-fixed to the laboratory (target-fixed) frame.

Bearing in mind the initial condition, the asymptotic time-independent hamiltonian in the initial channel is taken as $H_i^{as} = H_0 + V_{N1} + V_{N2} + V_{12}$, with $W_i(t) = V_{pHe}(t)$ being perturbation. And in the final channel we have $H_f^{as}(t) = H_0 + V_{p1}(t) + V_{N2}$ and $W_f(t) = V_{p2}(t) + V_{Np}(t) + V_{N1} + V_{12}$. The transition *prior*-amplitude in the time-dependent theory is

$$\tilde{a}_i^{fi}(\vec{b}) = -i \int_{-\infty}^{\infty} dt \langle \Phi^-(t) | W_i(t) | \Phi_0 \rangle \, e^{-i\varepsilon_0^{He} t}, \tag{13a}$$

while the *post*-amplitude is given by

$$a_f^{fi}(\vec{b}) = -i \int_{-\infty}^{\infty} dt \langle e^{i\hat{f}(\vec{v}_p,t)} \phi_{p1}, \varphi_{N2}^-(\vec{k}) | W_f(t) | \Phi^+(t) \rangle \, e^{i(\varepsilon^H + k^2/2)t}, \tag{13b}$$

where $e^{i\hat{f}(\vec{v}_p,t)}$ is the translational factor (see above). Both forms, (13a) and (13b), are equivalent, provided $\Phi^{\pm}(t)$ is the exact solution of Eq. (10) [15, 20]. Note that the authors of Refs. [10–12] unjustifiably set $W_f(t) = V_{12}$ in the amplitude (13b).

Comparing to Eq. (11), one readily sees the equivalence of the time-dependent and stationary formulations of scattering theory in the paraxial approximation. Finally, the relation between the T_{fi} and a_{fi} amplitudes is given by

$$T_{fi} = v_p \int d^2b \; e^{-i\vec{q}_\perp \cdot \vec{b}} a_{fi}(\vec{b}),$$

where $\vec{q} = \vec{p}_H - \vec{p}_0$, $\vec{q}_\perp = \vec{q} - q_z \vec{v}_p / v_p$, $\vec{v}_p = \{0, 0, v_p\}$, and $\vec{k} = \{k_\perp \cos\varphi_k, k_\perp \sin\varphi_k, k_z\}$. The q_z value is determined from energy conservation as $q_z = v_p/2 + Q/v_p$, where $Q = \varepsilon_0^{He} - \varepsilon_0^H - k^2/2$.

III. DISTORTED WAVE APPROXIMATIONS

From Eqs.(1), (2), (6), and (7) one can derive the higher Born terms as well as different versions of the distorted wave Born approximation (DWBA). They include, in particular, matrix elements which correspond to both Ne- and ee-Thomas mechanisms, where, however, all two-body potentials must be taken into account (see, for instance, different calculations presented in Ref. [2]). For example, in Ref. [2] the eikonal approximation was obtained, which introduces in the *prior*-form FBA matrix element (8b) a distorting phase factor,

$$|\phi_{p1}, \varphi_{N2}^-(\vec{k}), \vec{p}_H\rangle \to e^{(i/v_p)\hat{\delta}_f} |\phi_{p1}, \varphi_{N2}^-(\vec{k}), \vec{p}_H\rangle.$$

The details concerning its derivation on the basis of Eq. (11) in the closure approximation can be found in Ref. [2]. The result for the phase factor reads

$$\hat{\delta}_f = \ln\left(\frac{[v_p|\vec{R}(t) - \vec{x}| + \vec{v}_p \cdot (\vec{R}(t) - \vec{x})]^2 \; [v_p|\vec{R}(t) - \vec{r}_2| + \vec{v}_p \cdot (\vec{R}(t) - \vec{r}_2)]}{[v_p R(t) + \vec{v}_p \cdot \vec{R}(t)]^2 \; [v_p|\vec{R}(t) - \vec{x} - \vec{r}_2| + \vec{v}_p \cdot (\vec{R}(t) - \vec{x} - \vec{r}_2)]} \right).$$

We note that it is the asymptotic form of the product

$$e^{(i/v_p)\hat{\delta}_f} \to \Lambda_f^- = \Lambda_{p2}^- \Lambda_{pN}^- \Lambda_{N1}^- \Lambda_{12}^-, \tag{14}$$

with

$$\Lambda_{Z_1Z_2}^- = \exp\left(-\frac{\pi Z_1 Z_2}{2v_p}\right) \Gamma\left(1 - i\frac{Z_1Z_2}{v_p}\right) {}_1F_1\left[i\frac{Z_1Z_2}{v_p}, 1; -i(v_p r_{rel} + \vec{v}_p \vec{r}_{rel})\right].$$

Each factor $\Lambda_{Z_1Z_2}^-$ in (14) describes the distortion of interactions between different constituents of the two final compound systems, $p1$ and $N2$. In some sense, it is a 4C model (in analogy with well known 3C and 6C models in the scattering theory [22]).

The same procedure we can utilize in the case of the FBA matrix element in the *post*-form, replacing in (8a)

$$|\Phi_0, \vec{p}_0\rangle \to e^{(i/v_p)\hat{\delta}_i} |\Phi_0, \vec{p}_0\rangle.$$

Here again

$$e^{(i/v_p)\hat{\delta}_i} \to \Lambda_i^+ = \Lambda_{p1}^+ \Lambda_{p2}^+ \Lambda_{pN}^+, \tag{15}$$

and

$$\Lambda_{Z_1Z_2}^+ = \exp\left(-\frac{\pi Z_1 Z_2}{2v_p}\right) \Gamma\left(1 + i\frac{Z_1Z_2}{v_p}\right) {}_1F_1\left[-i\frac{Z_1Z_2}{v_p}, 1; i(v_p r_{rel} - \vec{v}_p \vec{r}_{rel})\right].$$

Each factor $\Lambda_{Z_1Z_2}^+$ in (15) describes the distortion due to interactions between the projectile proton and different constituents of the helium atom. This approximation is analogous to the 3C model.

Note that in the matrix elements we return to the time-independent formulation by replacing $v_p t \to R_z$. Both distortion factors, Λ_f^- and Λ_i^+, partially incorporate an effect of higher Born terms. However, they principally differ from those employed in the papers of Voitkiv *et al.*. The latter distortion factors clearly do not obey the correct boundary conditions $\Lambda_f^- \to 1$ and $\Lambda_i^+ \to 1$ at $t \to \pm\infty$, respectively, and lack justification from the multichannel scattering theory.

IV. NUMERICAL ILLUSTRATION

In Sec. II it was shown that in Refs. [10–12] the authors use the FBA matrix element in its *post*- form, leaving only the V_{12} contribution. Their account for distortion effects beyond the

FBA through the factors $\Lambda_{i,f}$ is, at least, questionable (see the above remarks). Moreover, in Ref. [12] the final distortion factor was neglected in calculations, $\Lambda_f = 1$, as being not so much significant. As remarked in Ref. [12], without the initial distortion factor, i.e., when $\Lambda_i = 1$, the contribution of the *ee*-Auger mechanism calculated there becomes much larger, while that of *ee*-Thomas vanishes. In view of these remarks, we might expect that neglecting the distortion effects in calculations does not reduce the role of the *ee*-Auger mechanism. Thus, in the FBA calculations presented in this section we set $\Lambda_i = \Lambda_f = 1$. Further, we use FBA in the *prior*-form (8b), pointing out once again that it is equivalent to the full, not truncated version of the *post*-form (8a). The quantity that was studied in Refs. [10–12] is the double differential cross section (DDCS)

$$\frac{d^2\sigma}{dk_\perp dk_z} = \frac{2k_\perp}{(2\pi)^5 v_p^2} \int\limits_0^{2\pi} d\varphi_k \int d^2q_\perp |T_{fi}|^2, \tag{16}$$

which describes a 2D distribution of the momentum components of the emitted electron ($k_x = k_\perp \cos\varphi_k,\ k_y = k_\perp \sin\varphi_k$).

Fig. 1 shows numerical results for the DDCS using the FBA amplitude in the *prior*-form (8b) with an accurate, highly correlated helium function of Mitroy *et al.* [24]. The kinematical situation is the same as that of Fig. 1 in Ref. [12]. However, while there is a common feature such as maximum located at $k_z = 0$, we find no maximum located at negative k_z values (approximately, at $k_z \approx -3.0$), in contrast to the results presented in Fig. 1 of Ref. [12]. The latter feature is, according to Voitkiv *et al.*, a clear signature of the *ee*-Auger mechanism. Thus, our numerical calculations that account for all the first-order (nonradiative) mechanisms using the accurate, highly correlated wave function of helium do not support the findings of Refs. [10–12]. In this connection, it should be noted that angular correlations due to the interaction V_{12} play very important role, if being included in the trial helium wave function $\Phi(\vec{r}_1, \vec{r}_2)$. They strongly influence the momentum distribution of the emitted electron in the backward direction, which is due to the shake-off mechanism [21]. And in Fig. 1 we can see manifestation of their effect in the region $k_z < 0$. However, it is quite different from manifestation of the *ee*-Auger mechanism claimed by Voitkiv *et al.* It is interesting that the *ee*-Auger mechanism has no manifestation in the OBK approximation either, as can be concluded from Fig. 2. According to Eq. (9), the effect of V_{12} found in Refs. [10–12] is clearly cancelled by the other first-order mechanism, which involves the target nucleus (the V_{N1} term). This finding markedly illustrates the importance of accounting for all the binary interactions between the particles taking part in the reaction.

V. SUMMARY AND CONCLUSIONS

We considered theoretically a transfer ionization channel in a fast proton-helium collision, focusing on the so-called first-order electron-electron capture mechanism proposed recently by Voitkiv *et al.* [10–12]. It was shown, using consistent quantum scattering theory, that this specific mechanism is included in a well known OBK scenario. The first-order calculations with an accurate trial helium function exhibited no signature of any significant effect of the *ee*-Auger process.

Some comments should be made with regard to the above theoretical analysis. The equivalence of the *post*- and *prior*-forms of the transition amplitude is realized only if the exact helium wave function is employed. In that case, we have the following equation for this function:

$$(\varepsilon_0^{He} - h_{10} - h_{20} - V_{N2})|\Phi_0\rangle = (V_{N1} + V_{12})|\Phi_0\rangle. \tag{17}$$

It can be readily shown, using energy conservation and properties of the final asymptotic state, that the projectile-electron potential V_{p1} in the left-hand side of Eq. (9) can be replaced with the operator $(\varepsilon_0^{He} - h_{10} - h_{20} - V_{N2})$ occurring in the left-hand side of Eq. (17). Thus, the left- and right-hand sides of the Schrödinger equation (17) correspond to, respectively, the *prior* and *post* matrix elements in Eq. (9). This feature explains a well documented fact that Eq. (9) is fulfilled to a good approximation in the case of an accurate trial helium function, which is typically obtained from a variational procedure. But if the trial function is poor, then the right-hand side of Eq. (17), which is related to the *post* matrix element in Eq. (9), appears to yield a better approximation to the exact result than in the case of the left-hand side. This observation explains why the results of Voitkiv *et al.* using the correlated and uncorrelated helium functions in their *post*-amplitude are similar (see Fig. 7 of Ref. [12]).

Finally, in their calculations Voitkiv *et al.* [10–12] adopt an approach in which they incoherently sum contributions to DDCS from different mechanisms, including, in particular, the OBK (or capture-shakeoff) and *ee*-Auger. However, such an approach is manifestly incorrect because, as follows from the above theoretical analysis, the *ee*-Auger contribution is already included in the OBK approximation. Thus, the corresponding numerical results of Refs. [10–12] can be regarded as erroneous.

Acknowledgments

This work is partially supported by the Russian Foundation for Basic Research (Grant No. 11-01-00523-a). Calculations are performed using the Moscow State University Research Computing Center (supercomputer Lomonosov). The authors are grateful to O. Chuluunbaatar and S. Houamer for their help. We are also grateful to R. Dörner and M. Schöffler for inspiring discussions.

---

[1] M. S. Schöffler *et al.*, Phys. Rev. A **79**, 064701 (2009); **80**, 042702 (2009).

[2] H.-K. Kim *et al.*, Phys. Rev. A **85**, 022707 (2012).

[3] V. Mergel *et al.*, Phys. Rev. Lett. **74**, 2200 (1995); D. Fischer *et al.*, J. Phys. B **35**, 1369 (2002).

[4] M. Schöffler *et al.*, J. Phys. B: At. Mol. Opt. Phys. **38**, L123 (2005); A. L. Godunov *et al.*, Phys. Rev. A **71**, 052712 (2005); M. S. Schöffler *et al.*, Phys. Rev. A **87**, 032715 (2013).

[5] L. H. Thomas, Proc. Roy. Soc. A **114**, 561 (1927).

[6] J. Eichler and T. Stöhlker, Physics Reports **439**, 1 (2007).

[7] J. R. Oppenheimer, Phys. Rev. **31**, 349 (1928); H. C. Brinkman and H. A. Kramers, Proc. Acad. Sci. Amsterdam **33**, 973 (1930).

[8] R. Shakeshaft, J. Phys. B: At. Mol. Phys. **7**, 1059 (1974)

[9] J. S. Briggs and K. Taulbjerg, J. Phys. B **15**, 2565 (1979).

[10] A. B. Voitkiv, B. Najjari, and J. Ullrich, Phys. Rev. Lett. **101**, 223201 (2008).

[11] A. B. Voitkiv, J. Phys. B **41**, 195201 (2008).

[12] A. B. Voitkiv and X. Ma, Phys. Rev. A **86**, 012709 (2012).

[13] J. S. Briggs and K. Dettmann, Phys. Rev. Lett. **33**, 1123 (1974).

[14] V. L. Shablov *et al.*, Phys. Part. Nucl. **41**, 335 (2010).

[15] J. R. Taylor, *Scattering Theory: the Quantum Theory of Nonrelativistic Collisions* (Wiley, New York, 1972).

[16] J. Dollard, J. Math. Phys. **5**, 729 (1964).

[17] E. Prugovečki E. and J. Zorbas, J. Math. Phys. **14**, 1358 (1973); Nucl. Phys. A. **213**, 541 (1973).

[18] R. K. Peterkop, *Theory of Ionization of Atoms* (Colorado Associated University Press, Boulder, 1977).

[19] J. M. Cheshire, Proc. Phys. Soc. **84**, 89 (1964).

[20] Dž. Belkić, R. Gayet, and A. Salin, Phys. Rep. **56**, 281 (1979).

[21] M. S. Schöffler *et al.*, Phys. Rev. A **87**, 032715 (2013).

[22] J. Berakdar, A. Lahmam-Bennani, and C. Dal Cappello, Phys. Rep. **374**, 91 (2003).

[23] S. Houamer, Yu. V. Popov, and C. Dal Cappello, Phys. Rev. A **81**, 032703 (2010).

[24] J. Mitroy, I. E. McCarthy, and E. Weigold, J. Phys. B **18**, 4149 (1985).



Figure 1: FBA distribution of the electron momentum components at $E_p = 3.6\,\mathrm{MeV}$. DDCS is given in barn/(a.u.)^2



Figure 2: The same as in Fig. 1, but in the OBK approximation.